

QualiPlate[™] Kit for LibertyLink[®] PAT/bar

Catalog Number AP 013

Highlights:

- High Sensitivity Protocol detects the PAT enzyme from the bar gene (PAT/bar) found in 0.1% StarLink corn
- Rapid Protocol screens individual seeds or leaf samples for the presence of PAT/bar

Contents of Kit:

- 1 antibody-coated 96-well plate
- PAT/bar Enzyme Conjugate
- 1 packet of Buffer Salts
- Substrate
- Stop Solution

Intended Use

The QualiPlate Kit for LibertyLink PAT/bar is designed for the qualitative laboratory detection of phosphinothricin acetyl transferase enzyme (PAT) coded for by the bar gene in grain, leaf, or seed. Two assay protocols are presented: The High Sensitivity Protocol will detect the PAT enzyme from the bar gene (PAT/bar) found in 0.1% StarLink® corn (by weight) and requires 2.5 hours to run. The Rapid Protocol (one hour total) is intended for use in screening individual seeds or leaf samples for the presence of PAT/bar. LibertyLink PAT from the pat gene, at concentrations present in T25 corn, is not detected in either format.

How the Test Works

This QualiPlate Kit is a "sandwich" Enzyme-Linked ImmunoSorbent Assay (ELISA). In the test, **corn** or **cotton** sample extracts are added to test wells coated with antibodies raised against PAT from the *bar* gene. Any residues present in the sample extract bind to the antibodies, and are then detected by addition of enzyme (horseradish peroxidase)-labeled PAT/*bar* antibody.

After a simple wash step, the results of the assay are visualized with a color development step; color development is proportional to PAT/bar concentration in the sample extract.

Lighter color = Lower concentration Darker color = Higher concentration

How the Kit Performs

The High Sensitivity Protocol will detect 0.1% StarLink corn grain. Samples are judged to contain more or less PAT/bar enzyme than 0.1% StarLink standard by comparing sample absorbances to the absorbance of a 0.1% StarLink sample extract (prepared by the user).

The Rapid Protocol is a strictly qualitative test, in which seed and leaf samples are screened for the presence or absence of the PAT/bar enzyme at the levels commonly seen in commercially modified crops. It is recommended that the user prepare extracts from known conventional and known PAT/bar expressing plant tissues to run in each assay as negative and positive controls.

Materials not Provided

- distilled or deionized water for preparing Wash/Extraction Buffer
- glass bottles or flask plus graduated cylinder with 1 liter capacity for preparation and storage of Wash/Extraction Buffer
- test or centrifuge tubes for extraction of grain
- Tissue Extraction Kit: snap-cap tubes and pestles for extraction of leaf samples (EnviroLogix Cat. No. ACC 002, 100/package)
- centrifuge capable of 5000 x g
- disposable tip, adjustable air-displacement pipettes which will measure 50 and 100 microliters (μL)
- marking pen (indelible)
- tape or Parafilm®



Prepare
Wash/Extraction Buffer

Preparation of Solutions

Wash/Extraction Buffer: Add the contents of the packet of Buffer Salts (phosphate buffered saline, pH 7.4 - Tween 20) to 1 liter of distilled or deionized water, and stir to dissolve. Store refrigerated when not in use; warm to room temperature prior to assay. If more extraction buffer is needed, order item # P-3563 from Sigma Chemical Co. (St. Louis, MO), or prepare the equivalent.

- timer
- microtiter ELISA plate reader
- wash bottle, or microtiter plate or strip washer
- multi-channel pipette that will measure 50 and 100 μL
- racked dilution tubes for loading samples into the plate with a multi-channel pipette, or the equivalent
- orbital plate shaker (optional)

Sample Preparation

Note: PAT/*bar* protein is not stable in solution. Extraction should be performed in 30 minutes or less. Do not extract overnight.

<u>High Sensitivity</u> Protocol for Ground Grain/Seed

This protocol requires that a small sample be analyzed. It is essential that this sample be well mixed and representative of the larger bulk. The test will detect 0.1% StarLink corn, containing PAT/bar (one kernel in a sample of 1000 kernels).

Note: Thorough mixing of the bulk grain sample and determination of an appropriate sampling plan are critical to the results of this testing, and are the responsibility of the user of this test kit. The USDA/GIPSA has prepared several guidance documents to address the issues involved in obtaining representative grain samples from static lots - such as trucks, barges, and railcars - and for taking samples from grain streams. Sampling plans should be chosen that best meet the needs of both the buyer and seller in terms of acceptable risks. Increasing the number of kernels in the sample and taking multiple samples will increase the likelihood of obtaining representative samples, and maximize the probability of detecting any contamination in the grain lot. For further information on USDA/GIPSA guidelines for obtaining representative samples and assessing detection probabilities for biotech grain, see the websites listed on page 3.

It is the responsibility of the user to ensure proper sampling and thorough mixing prior to analysis. Once representative samples have been obtained from the truck or container, they can be reduced in size using a splitter and uniformly ground and mixed. The finer the grind, the faster and more efficient the extraction.

- 1. Weigh 5 grams of ground corn sample into a 15 mL capacity vial or tube.
- 2. Add 12.5 mL of Wash/Extraction Buffer to each 5 gram sample. For all other grain sample sizes, add Wash/Extraction Buffer at the rate of 2.5 mL per gram of grain. Cap and shake vigorously by hand or vortex for 20-30 seconds. Let stand at room temperature to extract.
- 3. The extracted samples/controls must be clarified by centrifuging the extract at 5000 x g for 5 minutes. Insert a pipette tip below any floating lipid layer and above the pellet to remove the clarified sample.

Rapid Protocol for Screening Single Seeds or Leaf Punches:

Individual seeds:

- 1. Crush seeds: Seeds may be placed in a baggie and smashed with a hammer, then transferred to a tube; or, a seed crusher/48-well plate combination may be used (for example Hypure #HSC-100, PerkinElmer, Norton, OH, with Costar plate #3548, Corning Life Sciences, Acton, MA, or equivalent). Check to be sure that all seeds have been crushed. Take extreme care not to crosscontaminate between seed samples.
- Add 0.5 mL of Wash/Extraction Buffer to each crushed corn seed, or 1 mL to each crushed cotton seed. Mix for at least 30 seconds, then allow particles to settle.

Reference Websites:

www.gipsa.usda.gov/fgis/ handbook/gihbk1_inspec.aspx

USDA *Grain Inspection Handbook*, *Book 1*, *Grain Sampling*. This document provides a comprehensive overview of recommended sampling guidelines for static lots and grain streams. It reviews the various types of equipment and strategies that can be used to obtain a representative grain sample from different types of containers.

www.gipsa.usda.gov/fgis/biotech/sample2.htm

Guidance document entitled Sampling for the Detection of Biotech Grains, which provides important statistical sampling considerations when testing for the presence of biotech grains. It covers the basis for making probability determinations in accepting lots based upon different assumptions with respect to sample size, number of samples, sample preparation, etc.

www.gipsa.usda.gov/fgis/biotech/sample1.htm

Practical Application of Sampling for the Detection of Biotech Grains. This one-page application guide provides a table that gives sample sizes for selected lot concentrations and probability of rejecting the specified concentrations. It also provides a formula for making the calculation for other combinations.

www.gipsa.usda.gov/fgis/biotech/samplingplan1.xls This website provides a simple to use Sample Planner (29k Excel Spreadsheet). The planner allows you to enter different assumptions in terms of sample size, number of samples, acceptable quality level and to determine the probability of accepting lots with given concentration levels. It also plots the probabilities in graph form for easy interpretation. Specific data can be saved for documentation and future analyses.

Leaf testing:

- 1. Take a single leaf punch of approximately 5 millimeters diameter, using a micro-tube cap or a paper punch. Mash the leaf tissue with a pestle matched to the micro-tube, or with a disposable pipette tip, or a Hypure cutter (HCT-200, PerkinElmer, Norton, OH) in a 96-well plate (Costar #3370, Corning Life Sciences, Acton, MA, or equivalent).
- Add 0.25 mL of Wash/Extraction Buffer per leaf punch. Mix for at least 30 seconds, then allow particles to settle. Take extreme care not to cross-contaminate between leaf samples.

NOTE: It is recommended that the kit user extract known conventional and known PAT/*bar*-containing samples of the matrix being tested, and run these as Negative and Positive Controls (NC and PC) in each assay.

How to Run the Assay

- Read all of these instructions before running the kit.
- Allow all reagents to reach room temperature before beginning (at least 30 minutes with un-boxed plates and reagents at room temperature do not remove plates from bag with desiccant until they have warmed up).
- Organize all Control and sample extracts, and pipettes so that Step 1 can be performed in 15 minutes or less. The use of a multi-channel pipette is strongly recommended.
- Use a disposable-tip air-displacement pipette and a clean pipette tip to add each Calibrator and sample extract to the wells. Conjugate, Substrate, and Stop Solution may be added in the same manner; alternatively, use a repeating pipette with a disposable tip on the end of the Combitip for these three reagents.
- Use the well identification markings on the plate frame to guide you when adding the samples and reagents. In a qualitative assay, the Blank (BL), Negative Control (NC), and the Positive Control (PC) and 90 sample extracts (S) may be run in single wells on one plate. (See the Qualitative Assay Example Plate Layout Figure 1A).

HIGH SENSITIVITY PROTOCOL: The High Sensitivity Protocol will detect 0.1% StarLink corn (by weight) in ground grain/seed, and requires 2.5 hours of total assay incubation time.

Procedure

1. Add 50 μ L of PAT/bar-enzyme Conjugate to each well, followed immediately by 50 μ L of Wash/Extraction Buffer Blank (BL), 50 μ L of the centrifuged 0.1% StarLink ground corn and Negative ground corn extracts (PC and NC), and 50 μ L of each centrifuged sample extract (S) to their respective wells, as shown in the Example Plate Layout (Figure 1A)

NOTE: In order to minimize setup time it is recommended that a multi-channel pipette be used in steps 1, 5, and 7.

- 2. Thoroughly mix the contents of the wells by moving the plate in a rapid circular motion on the benchtop for a full 20-30 seconds. Be careful not to spill the contents!
- 3. Cover the wells with tape or Parafilm to prevent evaporation and incubate at ambient temperature for 2 hours. If an orbital plate shaker is available, shake plate at 200 rpm.
- After incubation, carefully remove the covering and vigorously shake the contents of the wells into a sink or other suitable container. Flood the wells



Allow all reagents to reach room temperature before beginning



Leaf punch



Extract sample



Centrifuge to clarify sample extract (High Sensitivity protocol only)

completely with Wash/Extraction Buffer, then shake to empty. Repeat this wash step three times. Alternatively, perform these four washes (300 $\mu L/\text{well})$ with a microtiter plate or strip washer. Slap the inverted plate on a paper towel to remove as much liquid as possible.

- 5. Add 100 μL of Substrate to each well.
- 6. Thoroughly mix the contents of the wells, as in step 2. Cover the wells with new tape or Parafilm and incubate for 30 minutes at ambient temperature. Use orbital shaker if available.
- 7. Caution: Stop Solution is 1.0N Hydrochloric acid. Handle carefully.
- 8. Add 100 μL of Stop Solution to each well and mix thoroughly. This will turn the well contents yellow.

NOTE: Read the plate within 30 minutes of the addition of Stop Solution.

RAPID PROTOCOL: The Rapid Protocol is less sensitive (able to detect 1% StarLink corn) but only requires one hour of total assay incubation time.

Procedure

1. Add 50 μL of PAT/bar-enzyme Conjugate to each well, followed immediately by 50 μL of Wash/Extraction Buffer Blank (BL), 50 μL of the user-prepared Positive and Negative Control extracts (PC and NC), and 50 μL of each sample extract (S) to their respective wells, as shown in the Example Plate Layout (Figure 1A).


NOTE: In order to minimize setup time it is recommended that a multi-channel pipette be used in steps 1, 5, and 7.

- 2. Thoroughly mix the contents of the wells by moving the plate in a rapid circular motion on the benchtop for a full 20-30 seconds. Be careful not to spill the contents!
- 3. Cover the wells with tape or Parafilm to prevent evaporation and incubate at ambient temperature for 45 minutes. If an orbital plate shaker is available, shake plate at 200 rpm.
- 4. After incubation, carefully remove the covering and vigorously shake the contents of the wells into a sink or other suitable container. Flood the wells completely with Wash/Extraction Buffer, then shake to empty. Repeat this wash step three times. Alternatively, perform these four washes (300 $\mu L/\text{well})$ with a microtiter plate or strip washer. Slap the inverted plate on a paper towel to remove as much liquid as possible.
- 5. Add 100 µL of Substrate to each well.
- 6. Thoroughly mix the contents of the wells, as in step 2. Cover the wells with new tape or Parafilm and incubate for 15 minutes at ambient temperature. Use orbital shaker if available.

Caution: Stop Solution is 1.0N Hydrochloric acid. Handle carefully.

7. Add 100 μ L of Stop Solution to each well and mix thoroughly. This will turn the well contents yellow.

NOTE: Read the plate within 30 minutes of the addition of Stop Solution.



Add Conjugate, controls and samples



Mix plate



Incubate



Wash plate



Read plates in a Plate Reader within 30 minutes of the addition of Stop Solution

How to Interpret the Results

Spectrophotometric Measurement

- 1. Set the wavelength of the microtiter plate reader to 450 nanometers (nm). (If it has dual wavelength capability, use 600, 630 or 650 nm as the reference wavelength.)
- 2. Set the plate reader to blank on the Wash/Extraction Buffer Blank wells (this should automatically subtract the mean optical density (OD) of the Blank wells from each control and sample OD). If the reader cannot do this, it must be done manually.

Note: The mean OD of the BLANK wells should not exceed 0.2.

Interpret the Results

For leaf and seed screening, compare the OD's of sample extracts to OD's of extracts from known leaf or seed samples (conventional and PAT/bar-expressing varieties). Note: very low positive results in a seed-screening assay may be due to cross-contamination during crushing. Truly positive seeds will result in OD readings typically greater than 0.500.

Figure 1A. Example of a typical Qualitative assay setup.

	1	2	3	4	5	6	7	8	9	10	11	12
Α	BL	S 6	S14	S22	S30	S38	S46	S54	S62	S70	S78	S86
В	NC	S 7	S15	S23	S31	S39	S47	S55	S63	S71	S79	S87
С	PC	S 8	S16	S24	S32	S40	S48	S56	S64	S72	S80	S88
D	S1	S9	S17	S25	S33	S41	S49	S57	S65	S73	S81	S89
Е	S2	S10	S18	S26	S34	S42	S50	S58	S66	S74	S82	S90
F	S3	S11	S19	S27	S35	S43	S51	S59	S67	S75	S83	BL
G	S4	S12	S20	S28	S36	S44	S52	S60	S68	S76	S84	NC
Н	S5	S13	S21	S29	S37	S45	S53	S61	S69	S77	S85	PC

Precautions and Notes

- Store all Plate Kit components at 4°C to 8°C (39°F to 46°F) when not in use.
- Do not expose Plate Kit components to temperatures greater than 37°C (99°F) or less than 2°C (36°F).
- Allow all reagents to reach ambient temperature (18°C to 27°C or 64°F to 81°F) before use.
- Do not use kit components after the expiration date.
- Do not use reagents or plates from one Kit with reagents or plates from a different Kit.
- Do not expose Substrate to sunlight during pipetting or while incubating in test wells.
- The assay has been optimized to be used with the protocol provided in the kit. Deviation from this protocol may invalidate the results of the test.
- As with all tests, it is recommended that results be confirmed by an alternate method when necessary.
- Observe any applicable regulations when disposing of samples and kit reagents.
- Use extreme caution to prevent sample-to-sample cross-contamination with samples, fluids, or disposables.



For Technical Support Contact Us At:

EnviroLogix 500 Riverside Industrial Parkway Portland, ME 04103-1486 USA

Tel: (207) 797-0300 Toll Free: 866-408-4597 Fax: (207) 797-7533

e-mail: info@envirologix.com

website: www.envirologix.com



LIMITED WARRANTY

EnviroLogix Inc. ("EnviroLogix") warrants the products sold hereunder ("the Products") against defects in materials and workmanship when used in accordance with the applicable instructions for a period not to extend beyond a product's printed expiration date. If the Products do not conform to this Limited Warranty and the customer notifies EnviroLogix in writing of such defects during the warranty period, including an offer by the customer to return the Products to EnviroLogix for evaluation, EnviroLogix will repair or replace, at its option, any product or part thereof that proves defective in materials or workmanship within the warranty period.

ENVIROLOGIX MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO ANY IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. The warranty provided herein and the data, specifications and descriptions of EnviroLogix products appearing in EnviroLogix published catalogues and product literature are EnviroLogix' sole representations concerning the Products and warranty. No other statements or representations, written or oral, by EnviroLogix' employees, agents or representatives, except written statements signed by a duly authorized officer of EnviroLogix Inc., are authorized; they should not be relied upon by the customer and are not a part of the contract of sale or of this warranty.


EnviroLogix does not warrant against damages or defects arising in shipping or handling, or out of accident or improper or abnormal use of the Products; against defects in products or components not manufactured by EnviroLogix, or against damages resulting from such non-EnviroLogix made products or components. EnviroLogix passes on to customer the warranty it received (if any) from the maker thereof of such non-EnviroLogix made products or components. This warranty also does not apply to Products to which changes or modifications have been made or attempted by persons other than pursuant to written authorization by EnviroLogix.


THIS WARRANTY IS EXCLUSIVE. The sole and exclusive obligation of EnviroLogix shall be to repair or replace the defective Products in the manner and for the period provided above. EnviroLogix shall not have any other obligation with respect to the Products or any part thereof, whether based on contract, tort, strict liability or otherwise. Under no circumstances, whether based on this Limited Warranty or otherwise, shall EnviroLogix be liable for incidental, special, or consequential damages.


This Limited Warranty states the entire obligation of EnviroLogix with respect to the Products. If any part of this Limited Warranty is determined to be void or illegal, the remainder shall remain in full force and effect.


Parafilm is a registered trademark of American Can Corporation LibertyLink is a trademark of Bayer StarLink is a trademark of Aventis CropScience EnviroLogix, the EnviroLogix logo, and QualiPlate are trademarks of EnviroLogix Inc.

© EnviroLogix 2015











SEC	TION 3. Composi	ion/inform	ation on ing	redients				
3.2	Mixture Aqueous solution	IN Hydroch	lorie Acid (1)	N HCl, 3 % HCl)				
	Chemical name	Amount (%)	CAS No	Classification According to OSI	IA 29CFR 1910.1200			
	Hydrochloric acid	1-4%	EC No	Hazard Classification	Hazard Code			
	TIJ MECHANIS	***************************************	7647-01-0					
			231-595-7	May be Corrosive to Metals	H290			
				Causes Skin Irritation	H315			
				Causes Serious Eye Damage	H318			
room	THAN 1 P							
	THON 4. First aid Description of first ai							
A	After inhalation :	i iiwasures		In case of inhalation. Remove to fresh air. I	f not breathing give artificial			
After skin contact :				respiration. Get medical attention immediately. In case of skin contact, Remove contaminated clothing and shoes immediately. Wash affected area with mild scop or detergent for at least 10 minutes or until r				
A	After eye contact :			evidence of chemical remains. In case of eve contact, immediately flush eye minutes. Lifting eyelids occasionally, until r medical attention immediately.	es with plenty of water for at leas no evidence of chemical remains			
A	After swallowing:			In case of ingestion, DO NOT Induce vomit medical personnel. Never give anything by a physician immediately.				
4.2 N	dost important symp And delayed:	toms and ef	lects, both acr	May cause skin irritation and eye damage				
	ndication of any imm pecial treatment nee		cal attention	and DO NOT use sodium bicarbonate in an atten	apt to neutralize the acid.			
SEC	TION 5. Firefight	ng measur	res.					
	xtinguishing media:	ng measur		CO2, extinguishing powder or water spray. Fight	arger fires with water spray or a			
575	special hazards arisir	a from the	mbetance or	resistant foam.				
	nixture:	g mountine	substance or	Hydrogen Chloride gas				
5.3 Advice for firefighters:				Wear protective gear appropriate for fire conditions including respiratory protective gear.				
6.1 F	CTION 6. Accident Personal precautions, and emergency process	protective of		In the case of spilled mixture wear gloves to prever spill, additional protection is recommended	at skin contact. In the case of a l			
	Environmental preca	attenne.		Do not discharge mixture to sewer system or water	*****			
	anvironmentai preca	itions:		DO HOT describings illivings to sewer system of water	mayo.			
6.2 I								

Practice good che clothing. Store in tightly cl- sunlight and heat.	emical hygiene when handling. A cosed, non-metal container, in a c Store in well aired storage room	other specific uses are stipulated USA (OSHA)
Store in tightly el- sunlight and heat. Apart from the us tection Hydrogen	osed, nen-metal container; in a c Store in well aired storage room ses mentioned in section 1.2, no. European (Commission directive 96/94) BHr TWA = 5 ppm	comosive compatible area. Prevent directs other specific uses are stipulated USA (ONIA)
Store in tightly el- sunlight and heat. Apart from the us tection Hydrogen	osed, nen-metal container; in a c Store in well aired storage room ses mentioned in section 1.2, no. European (Commission directive 96/94) BHr TWA = 5 ppm	comosive compatible area. Prevent directs other specific uses are stipulated USA (ONIA)
Store in tightly el- sunlight and heat. Apart from the us tection Hydrogen	osed, nen-metal container; in a c Store in well aired storage room ses mentioned in section 1.2, no. European (Commission directive 96/94) BHr TWA = 5 ppm	comosive compatible area. Prevent directs other specific uses are stipulated USA (ONIA)
Apart from the us tection Hydrogen	Store in well aired storage roomses mentioned in section 1.2, no- ses mentioned in section 1.2, no- section	ns. other specific uses are stipulated USA (OSHA)
(ection	European (Commission directive 96/94) 8Hr TWA = 5 ppm	USA (OSHA)
Hydrogen	directive 96/94) 8Hr TWA = 5 ppm	
Hydrogen	directive 96/94) 8Hr TWA = 5 ppm	
	directive 96/94) 8Hr TWA = 5 ppm	
	(v > mk/m2)	Ceiling Limit = 5 ppm (7.5 mg/m3)
	STEL = 10 ppm (15 mg/m3)	
	V II	At .
The usual precaut	tionary measures should be adhe	ered to when handling chemicals.
approved under a Eye and face prot	ppropriate government standard tection regulations are described	s such as NIOSH (US) or EN 166 (EU) by OSHA (US) in 29CFR1910.133. Do
technique (withou Dispose of contan laboratory practic	ut touching glove's outer surface minated gloves after use in accor- ces. Wash and dry hands. The sel	 to avoid skin contact with this product dance with applicable laws and good lected protective gloves have to satisfy
using risk analysi may be used as a	s protocols. An approved dispos backup to engineering controls.	able air purifying particulate respirator Always use respirators and components
Contain spills, do	not allow into environment	
	Penision - 13 April 2015	Page 3
EC CETE PETER	general or local e exposure limits. The usual precau Safety glasses wi approved under a Exposure of the pro- tot wear centact. Handle with glow technique (without Dispose of contait aboratory practic the specifications. Appropriate responsating in the suiting risk analysis may be used as a tested and approx (EU).	Facilities using this moture should be equipped separate or local enhant verification to keep article exposure laints. The usual precurationary measures schoold be admits shortly glauses with side shields, avagles. Use or gappe of leader appropriate powerment standary groups and south and state of the

9.1 Information on basic physical and				
chemical properties:				
a) Appearance:	Clear liquid, colorless to slight	wallow		
a) Appearance: b) Odor:	Pungent (slight)	yanow.		
e) Odor Threshold	No Data Available			
d) pH:	pH 1			
	No Data Available			
e) Melting point/freezing point:	No Data Available.			
f) Boiling point/Boiling range:	No Data Available. Not applicable.			
g) Flash point:		Part Control 1		
h) Evaporation rate:	0.36 (Water) compared with n	-Butyl Acetate = 1		
i) Flammability (solid, gaseous):	No Data Available			
j) Upper/lower flammability or explosive limits:	No Data Available			
k) Vapor pressure:	No Data Available			
Vapor pressure: Vapor density	No Data Available			
m) Relative density:	No Data Available			
n) Solubility(ies):	Fully miscible, water.			
Solubility(ies): O Partition Coefficient: n-Octanol/water:	No Data Available			
p) Auto-ignition temperature:	No Data Available			
Auto-ignition temperature: Decomposition temperature:	No Data Available			
		Englanding to the America	10.00	
r) Viscosity:	No Data Available but should	be sitting to that of v	vater	
s) Explosive properties:	No Data Available.			
t) Oxidizing properties:	No Data Available	4.44		
9.2 Other information.	No further relevant information	n available.		
ECTION 10. Stability and reactivity				
10.1 Reactivity:	No data available			
10.2 Chemical Stability:	Stable under normal tempera	tures and pressures		
and the state of t	statens in server \$1.00	. 8		
10.3 Possibility of hazardous reactions:	Under normal conditions of	storage and use, naza	ruous reactions will	not occur.
10 4 Conditions to avoid:	No specific data			
10.5 Incompatible materials:	Metals, Alkali metals, bases,	Amines.		
10.6 Hazardous decomposition products:	Under normal conditions of			
	not be produced.	dorage and use, haza	rdous decomposition	ns products should
SECTION 11. Toxicological information	not be produced.			
Information on Toxicological Effects	not be produced.	Effect Dose	Specia	es
Information on Toxicological Effects	not be produced. 7647-01-0 HC1 Acute oral toxicity	Effect Dose	Specia	es
Information on Toxicological Effects	not be produced. 7647-01-0 HC1 Acute oral toxicity Acute dermal toxicity	Effect Dose LD50=900mg/kg No data	Special rabbit	es
Information on Toxicological Effects	not be produced. 7647-01-0 HC1 Acute oral toxicity	Effect Dose	Special rabbit	es
Information on Toxicological Effects Acute effects (toxicity tests):	7647.01.0 HCI Acute oral toxicity Acute inhalative toxicity	Effect Dose LD50-900mg/kg No data LC50 = 3124 mg	Special rabbit	es
Information on Toxicological Effects Acute effects (toxicity tests): Sensitization:	7647.01-0 HCI Acute creal loxicity Acute dermal toxicity Acute including the content of the cont	Effect Dose LD50-900mg/kg No data LC50 = 3124 mg	Special rabbit	es
Information on Toxicological Effects Acute effects (toxicity tests). Sensitization: CMR (carcinogenicity, matagenicity and toxici	7647.01.0 HCI Acute oral toxicity Acute inhalative toxicity No sensitizing effects known by	Effect Dose LD50-900mg/kg No data LC50 = 3124 mg	Special rabbit	es
Information on Toxicological Effects Acute effects (toxicity tests): Semsitization: CMR (carcinogenicity, mutagenicity and toxicit for reproduction) effects:	net be produced. 7647-81-0-HCI Acute and towardy Acute demail socialy. Acute insulative towardy No sensitizing effects known y No CMR effects.	Effect Dose LD50-900mg/kg No data LC50 = 3124 mg	Special rabbit	es
Information on Toxicological Effects Acute effects (toxicity tests). Sensitization: CMR (carcinogenicity, matagenicity and toxici	7647.01.0 HCI Acute oral toxicity Acute inhalative toxicity No sensitizing effects known by	Effect Dose LD50-900mg/kg No data LC50 = 3124 mg	Special rabbit	es
Information on Toxicological Effects Acute effects (toxicity tests): Semsitization: CMR (carcinogenicity, mutagenicity and toxicit for reproduction) effects:	net be produced. 7647-81-0-HCI Acute and towardy Acute demail socialy. Acute insulative towardy No sensitizing effects known y No CMR effects.	Effect Dose LD50-900mg/kg No data LC50 = 3124 mg	Special rabbit	es
Information on Toxtoological Effects Acute effects (toxicity tests). Sensitization. CMR (carcinogenicity, mutagenicity and toxicit for reproduction) effects: Additional toxicological information:	net be produced. 7647-81-0-HCI Acute and towardy Acute demail socialy. Acute insulative towardy No sensitizing effects known y No CMR effects.	Effect Dose LD50-900mg/kg No data LC50 = 3124 mg	Special rabbit	es
Information on Toxtoological Effects Acute effects (toxicity tests): Sensitization: CMR (carcingenicity, mutagenicity and toxicity for reproduction) effects: Additional toxicological information: ECTION 12. Ecological information	net be produced. **GET-AL-B-HC!* Acute cond testorb. Acute dermal testorb. Acute dermal testorb. Acute inhalative testocity No sensitizing effects known by No CMR effects. No Additional Information	Effect Dose LD59-900mg/kg No data LC59 = 3124 mg	Special rabbit rat	es
Information on Toxtoological Effects Acute effects (toxicity tests): Sensitization: CMR (carcingenicity, mutagenicity and toxicity for reproduction) effects: Additional toxicological information: ECTION 12. Ecological information	net be produced. 7647-01-0 HCI Acute ceal toxicity Acute inhalter toxicity Acute inhalter toxicity No sensitizing effects known No CMR effects. No Additional Information	Effect Dose LD50=90mg/kg No data LC50 = 3124 mg	Special rubbit rat	Species
Information on Toxtoological Effects Acute effects (toxicity tests): Sensitization: CMR (carcingenicity, mutagenicity and toxicity for reproduction) effects: Additional toxicological information: ECTION 12. Ecological information	net be produced. 7647-01-0 HC1 Acute end nosicity Acute malastre tenesity Acute malastre tenesity No emissizing effects known No Additional Information Aquatic texicity (IN IRC) Acute file November (IN IRC) Acute file November (IN IRC)	Effect Dose LD59-900mg/k; No data LC59 = 3124 mg	Special rabbit rat	es
Information on Toxicological Effects Acute effects (toxicity tests); Sensitization: CMR (carcinogenicity, mutagenicity and toxicit for reproduction) effects;	net be produced. 7647-01-0 HCI Acute ceal toxicity Acute inhalter toxicity Acute inhalter toxicity No sensitizing effects known No CMR effects. No Additional Information	Effect Dose LD50=90mg/kg No data LC50 = 3124 mg	Special rubbit rat	Species